

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 260 501 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
27.11.2002 Bulletin 2002/48

(51) Int Cl.7: **C07C 319/20, C07C 323/60**

(21) Application number: **02008248.3**

(22) Date of filing: **18.04.2002**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

- Lexner, Jael
Tel Aviv 63403 (IL)
- Saeed, Shadi
Haifa 33266 (IL)
- Kaspi, Joseph
Givatayim 53201 (IL)
- Leman, Ori
Givatayim 53294 (IL)

(30) Priority: **13.05.2001 IL 14310601**


(71) Applicant: **CHEMAGIS LTD.**
Tel Aviv 61090 (IL)

(72) Inventors:
• Naddaka, Vladimir
Lod 71456 (IL)
• Menashe, Naim
Herzliya (IL)

(74) Representative:
**Forstmeier, Dietmar, Dr. rer. nat., Dipl.-Chem. et
al
Boeters & Bauer,
Berlineranger 15
81541 München (DE)**

(54) **Process for the preparation of diphenylmethylthioacetamide**

(57) The invention provides a process for the preparation of diphenylmethylthioacetamide (I) as described in Scheme 4 comprising reacting of the isothiuronium salt or its corresponding base of the formula IV with an acetamide of the formula XCH_2CONH_2 , wherein X represents a halogen, M represents an alkali metal and A represents an anion, in a protic medium at a temperature of less than 100° C.



Schem 4

EP 1 260 501 A1

Description

[0001] The present invention relates to an improved process for the preparation of diphenylmethylthioacetamide (I), a key intermediate in the synthetic pathway for Modafinil.

5 [0002] Modafinil is defined as a CNS stimulant and is marketed under the trade name "Provigil", for the treatment of narcolepsia.


Modafinil was introduced by Lafon. Its preparation, use and pharmacological properties were described in US Patent 4177290.

10 Prior Art

[0003] US Patent no. 4177290 describes the synthetic pathways for the preparation of Modafinil (III). Both of the pathways described therein use diphenylmethylthioacetic acid (II) as starting material.

15

20



25

II

[0004] One synthetic method is based on the reaction of diphenylmethylthioacetyl chloride (obtained from II and thionyl chloride) with ammonia, followed by oxidizing the thioamide formed with hydrogen peroxide. Total yield, as reported, is 63% (Scheme I).


30

35

40

45


50



Scheme I

55


[0005] A different synthetic pathway described in the same patent involves the oxidation of II as first stage, followed by esterification with dimethylsulfate and amidation with gaseous ammonia. The total yield is 41% (Scheme 2).



Scheme 2

25

[0006] Compound II is prepared by the reaction of diphenylmethanol with thiourea in the presence of HBr followed by basic hydrolysis and reaction with chloroacetic acid (Scheme 3).



Scheme 3

45

[0007] Both synthetic methods involve four chemical steps (starting from diphenylmethanol) and the yields, as reported, are moderate.

[0008] Furthermore, both pathways involve toxic and dangerous reagents such as dimethylsulfate, ammonia and thionyl chloride.

50

[0009] In order to improve the synthetic method, according to the present invention there is now reacted the isothiuronium salt (IV) (obtained from the reaction between diphenylmethanol and thiourea) with haloacetamide.

[0010] According to the present invention it has now been found that haloacetamide reacts with the isothiuronium salt (IV) in protic solvents (like alcohols or water) at temperatures ranging from 0 - 100° C in the presence of a strong base, such as alkali metal hydroxide.

55

[0011] Preferably, chloroacetamide is reacted with the said isothiuronium salt in water at 60-70° C using sodium hydroxide as base.

[0012] When the reaction is carried out under the above-mentioned conditions diphenylmethanesulfonylacetamide is obtained in 95% yield from diphenylmethanol.

[0013] Thus according to the present invention there is now provided a process for the preparation of diphenylmeth-ylthioacetamide (I) as described in Scheme 4 comprising reacting of the isothiuronium salt or its corresponding base of the formula IV with an acetamide of the formula XCH_2CONH_2 , wherein X represents a halogen, M represents an alkali metal and A represents an anion, in a protic medium at a temperature of less than 100° C.

5


10

15

20

25

30



Scheme 4

[0014] The said method comprises only three chemical steps. It does not involve toxic or corrosive reagents. Very pure Modafinil is obtained at the end of the synthetic pathway at 67% yield (starting from diphenylmethanol). Thus, diphenylmethanol is reacted with thiourea in the presence of hydrobromic acid in aqueous medium under reflux and then cooled. The isothiuronium salt (IV) formed is filtered and reacted with chloroacetamide under aqueous base conditions at 60-70° C. The solution is cooled and the diphenylmethylthioacetamide formed is filtered and collected.

[0015] All the physical properties and the NMR and IR spectra are in perfect agreement with the proposed structure.

[0016] The following examples will serve as further illustration and clarification of the present invention.

[0017] While the invention will now be described in connection with certain preferred embodiments in the following examples so that aspects thereof may be more fully understood and appreciated, it is not intended to limit the invention to these particular embodiments. On the contrary, it is intended to cover all alternatives, modifications and equivalents as may be included within the scope of the invention as defined by the appended claims. Thus, the following examples which include preferred embodiments will serve to illustrate the practice of this invention, it being understood that the particulars shown are by way of example and for purposes of illustrative discussion of preferred embodiments of the present invention only and are presented in the cause of providing what is believed to be the most useful and readily understood description of formulation procedures as well as of the principles and conceptual aspects of the invention.

Reference Example 1.

Preparation of isothiuronium salt (IV).

[0018] Diphenylmethanol (130 g, 0.7 mole) and thiourea (65 g, 0.85 mole) are added in 0.5 l reactor charging with water (325 ml). The mixture is heated to 95° C (an emulsion is obtained) and 48% HBr 260 gr. 3.22 mole, 4.6 equivalents) is then added gradually during 0.5 hour. The mixture is heated under reflux (106 - 107° C) for 0.5 hour and cooled to 80 - 85° C. At this temperature, the mixture is seeded with several crystals of the product and the mixture is stirred at that temperature for 0.5 hour and then cooled to 25°. The colorless crystals are collected by filtration, washed with water (200 ml) yielding about 240 gr. of wet crude isothiuronium salt.

Preparations of diphenylmethylthioacetamid

Example 1a.

5 [0019] A 2 L reactor was charged with diphenylmethylisothiuronium bromide crude wet obtained in reference example 1 (240 gr.) and water (700 ml.) under nitrogen. The suspension was heated to 60° C and 46% aqueous NaOH solution (98 ml, 1.68 mole, 2.4 eq.) was added. The reaction mixture was heated to 85° C and stirred until all the solid was dissolved. Then, it was cooled to 60° C and chloroacetamide (80 g, 0.84 mole, 1.2 eq.) was added in five portions
 10 hour at 60 - 70° C during one hour. The suspension is stirred at 70° C for 4-5 hours. The mixture was filtered while warm and the cake was washed with hot water (250 ml). Diphenylmethylthioacetamide crude wet is obtained [220 gr., HPLC assay: 78%, HPLC purity: 95%, yield: 95% (from diphenylmethanol.)]
 20 gr. of the product was recrystallized twice from ethyl acetate, dried *in vacuo* to give 15 gr. of pure title compound.
¹H-NMR (CDCl₃); δ(ppm) = 3.3 (2H, s, CH₂), 5.18 (1H, s, CH), 6.54 (2H, bs, NH₂), 7.18-7.43 (10H, m, 2 x Ph).
¹³C-NMR (CDCl₃); δ(ppm) = 35.55 (CH₂), 54.56 (CH), 127.60, 128.26, 140.20 (Ph), 171.82 (C=O).

Example 1b.

[0020] A 0.5 L reactor was charged with diphenylmethylisothiuronium bromide crude wet (61 gr.) and water (163 ml.) under nitrogen. The suspension was heated to 40° C and 46% aqueous NaOH solution (25 ml.) was added.
 20 The reaction mixture was heated to 80° C and stirred until all the solid was dissolved. Then, it was cooled to 30° C and chloroacetamide (19.8 g, 0.21 mole, 1.2 eq., purity 98%) was added in four portions during one hour. The suspension was stirred at 30 - 35° C for 4-5 hours. Then the mixture was filtered while warm and the cake was washed with water and a mixture of methanol and water (4:1). The product was dried on air to give diphenylmethylthioacetamide crude wet was obtained [78.3 gr., HPLC assay: 88%, HPLC purity: 88%, yield: 88% (from diphenylmethanol.)]

Example 1c.

[0021] A 0.5 L reactor was charged with diphenylmethylisothiuronium bromide crude wet (61 g.) and methanol (163 ml.) under nitrogen. The suspension was heated to 40 - 50° C and 46% aqueous NaOH solution (25 ml.) was added.
 30 The reaction mixture was heated to 60° C and stirred for ½ hour. Chloroacetamide (19.8 g, 0.21 mole, 1.2 eq., purity 98%) was added in two portions during 10 minutes. The suspension was stirred at heating under reflux for 3-4 hours. Then water (80 ml.) was added dropwise at 60 - 65° C. The mixture was cooled to 30° C and filtered. The cake was washed with a mixture of methanol and water (4:1) and water. Diphenylmethylthioacetamide crude wet was obtained [34.3 g, HPLC assay: 95%, HPLC purity: 99%, yield: 71% (from diphenylmethanol.)]

Example 1d.

[0022] A 0.5 L reactor was charged with diphenylmethylisothiuronium bromide crude wet (38 g.) and water (117 ml.) under nitrogen. The suspension was heated to 40 - 50° C and 46% aqueous NaOH solution (16.4 ml.) was added.
 40 The reaction mixture was heated to 80° C until all the solid had been dissolved. Then it was cooled to ca. 60° C and bromoacetamide (20 g, 0.142 mole, 1.2 eq., purity 98%) was added in five portions during one hour. The suspension was stirred at 70° C for 4-5 hours. Then it was filtered while warm and the cake washed with hot water. Diphenylmethylthioacetamide crude wet was obtained [32.7 g, HPLC assay: 78%, HPLC purity: 99%, yield: 90% (from diphenylmethanol.)]

Reference Example 2.


[0023] A 1.0 L reactor was charged with diphenylmethylthioacetamide crude wet (220 gr.) obtained from example 1a and glacial acetic acid (610 ml). The mixture was heated to 40° C and stirred until full dissolution is achieved. 5.8% H₂O₂ solution (500 g, 1.2 eq.) was added dropwise during 0.5 hours at 40 - 45° C. The reaction mixture was stirred at 40 - 45° C for 4 hours. Then sodium metabisulfite (18.3 gr.) in 610 ml water was added in order to quench the unreacted H₂O₂ and the suspension was stirred for 0.5 hours. Then the reaction mixture was cooled to 15° C and filtered. The cake was washed with water (610 ml.) and dried on air to obtain crude wet Modafinil (205 g). Reslurry in refluxing ethyl acetate, followed by recrystallization from methanol:water (4:1) solution afforded pure Modafinil [(125 g, HPLC assay: 99.9%, HPLC purity: 99.9%, yield: 67% (from diphenylmethanol.)]

[0024] It will be evident to those skilled in the art that the invention is not limited to the details of the foregoing illustrative examples and that the present invention may be embodied in other specific forms without departing from the essential attributes thereof. It is therefore desired that the present embodiments and examples be considered in

all respects as illustrative and not restrictive, reference being made to the appended claims, rather than to the foregoing description and all the changes which come within the meaning and range of equivalency of the claims are therefore intended to be embraced therein.

Claims

1. A process for the preparation of diphenylmethylthioacetamide (I) as described in Scheme 4 comprising reacting of the isothiuronium salt or its corresponding base of the formula IV with an acetamide of the formula XCH_2CONH_2 , wherein X represents a halogen, M represents an alkali metal and A represents an anion, in a protic medium at a temperature of less than 100° C.



Scheme 4

2. A process according to claim 1, where X is selected from the group consisting of chlorine and bromine atoms.
3. A process according to claims 1 or 2 where the reaction takes place at a temperature between 60 - 70° C.
4. A process according to claims 1, 2 or 3 where M is a sodium atom.
5. A process wherein diphenylmethylthioacetamide produced according to any one of the preceding claims is used for the preparation of Modafinil.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 02 00 8248

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	DATABASE CROSSFIRE BEILSTEIN 'Online! Beilstein Institut zur Förderung der Chemischen Wissenschaften, Frankfurt am Main, DE; Database accession no. Reaction ID 4315629 XP002211381 * abstract * & TIMMERMANN, H. ET AL.: ARZNEIM. FORSCHUNG; 20; 1970, 1258-59,	1-4	C07C319/20 C07C323/60
A,D	US 4 177 290 A (LAFON LOUIS) 4 December 1979 (1979-12-04) * examples 1,1A *	1-4	
X		5	
A	EP 0 097 071 A (LAFON LABOR) 28 December 1983 (1983-12-28) * preparation I *	1-4	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			C07C
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 28 August 2002	Examiner Seufert, G
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03 82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 00 8248

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

28-08-2002

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 4177290	A	04-12-1979	GB 1584462 A	11-02-1981
			BE 865468 A1	02-10-1978
			CA 1091679 A1	16-12-1980
			CH 628026 A5	15-02-1982
			DE 2809625 A1	05-10-1978
			DK 140878 A ,B,	01-10-1978
			ES 468378 A1	16-12-1978
			FR 2385693 A1	27-10-1978
			IE 46566 B1	27-07-1983
			IT 1094041 B	26-07-1985
			JP 1400453 C	28-09-1987
			JP 53121724 A	24-10-1978
			JP 62009103 B	26-02-1987
			LU 79335 A1	28-09-1978
			LU 90153 A9	16-02-1998
			NL 7803432 A ,C	03-10-1978
EP 0097071	A	28-12-1983	FR 2528038 A2	09-12-1983
			AT 15185 T	15-09-1985
			CA 1199916 A1	28-01-1986
			DE 3360668 D1	03-10-1985
			DK 253583 A ,B,	05-12-1983
			EP 0097071 A1	28-12-1983
			ES 522981 D0	16-03-1984
			ES 8403445 A1	16-06-1984
			IE 54988 B1	11-04-1990
			JP 59001459 A	06-01-1984

EPO FORM P0458

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82